



# NAIMA



## TABLE VARIETY

- *Excellent productivity.*
- *Medium early variety.*
- *Big oval to long oval tubers.*
- *White flesh.*

### PLANT & TUBER

FOLIAGE MATURITY	MEDIUM EARLY
EMERGENCE	MODERATE
PLANT VIGOUR	HIGH
SKIN COLOUR	YELLOW
FLESH COLOUR	WHITE
TUBER SHAPE	OVAL TO LONG OVAL
SHAPE UNIFORMITY	GOOD
EYE DEPTH	SHALLOW
TUBER SIZE	LARGE

### PESTS & DISEASES

RESISTANCE TO FOLIAGE LATE BLIGHT	FAIRLY LOW
RESISTANCE TO TUBER LATE BLIGHT	FAIRLY LOW
RESISTANCE TO PITTED COMMON SCAB	MODERATE
RESISTANCE TO NETTED COMMON SCAB	MODERATE
RESISTANCE TO PVY	FAIRLY LOW
RESISTANCE TO PCN <i>G.rostochiensis</i> (Ro1-4)	LOW
RESISTANCE TO BLACK WART	HIGH

### QUALITY & USE

YIELD	VERY HIGH
DRY MATTER CONTENT	LOW
COOKING TYPE	A
DISINTEGRATION	VERY SLIGHT
DARKENING AFTER COOKING	NONE
FRY COLOUR at harvest	MEDIUM
STORAGE ABILITY	FAIRLY HIGH

### PHYSIOLOGY

DORMANCY	MEDIUM
RESISTANCE TO EXTERNAL DAMAGE	FAIRLY HIGH
RESISTANCE TO INTERNAL DAMAGE	FAIRLY HIGH
RESISTANCE TO INTERNAL RUST SPOT	FAIRLY HIGH
RESISTANCE TO DESPROUTING	HIGH